

corresponds to a particular nucleotide sequence. In Figures 4-6, the principle of array-based hybridization assays is illustrated with respect to the single feature 404 to which a number of identical probes 405-409 are bound. In practice, each feature of the array contains a high density of such probes but, for the sake of clarity, only a subset of these are shown in Figures 4-6.

Once an array has been prepared, the array may be exposed to a sample solution of target DNA or RNA molecules (410-413 in Figure 4) labeled with fluorophores, chemiluminescent compounds, or radioactive atoms 415-418. Labeled target DNA or RNA hybridizes through base pairing interactions to the complementary probe DNA, synthesized on the surface of the array. Figure 5 shows a number of such target molecules 502-504 hybridized to complementary probes 505-507, which are in turn bound to the surface of the array 402. Targets, such as labeled DNA molecules 508 and 509, that do not contain nucleotide sequences complementary to any of the probes bound to array surface, do not hybridize to generate stable duplexes and, as a result, tend to remain in solution. The sample solution is then rinsed from the surface of the array, washing away any unbound labeled DNA molecules. Finally, the bound labeled DNA molecules are detected via optical or radiometric scanning. Figure 6 shows labeled target molecules emitting detectable fluorescence, radiation, or other detectable signal. Optical scanning involves exciting labels of bound labeled DNA molecules with electromagnetic radiation of appropriate frequency and detecting fluorescent emissions from the labels, or detecting light emitted from chemiluminescent labels. When radioisotope labels are employed, radiometric scanning can be used to detect the signal emitted from the hybridized features. Additional types of signals are also possible, including electrical signals generated by electrical properties of bound target molecules, magnetic properties of bound target molecules, and other such physical properties of bound target molecules that can produce a detectable signal. Optical, radiometric, or other types of scanning produce an analog or digital representation of the array as

10020333-1